



US005223263A

United States Patent [19]**Hostetler et al.**[11] **Patent Number:** **5,223,263**[45] **Date of Patent:** **Jun. 29, 1993**[54] **LIPONUCLEOTIDE-CONTAINING LIPOSOMES**[75] **Inventors:** **Karl Y. Hostetler, Del Mar; Raj Kumar, San Diego; Louise M. Stuhmiller, Rancho Santa Fe, all of Calif.**[73] **Assignee:** **Vical, Inc., San Diego, Calif.**[21] **Appl. No.:** **373,088**[22] **Filed:** **Jun. 28, 1989****Related U.S. Application Data**

[63] Continuation-in-part of Ser. No. 216,412, Jul. 7, 1988, abandoned, and a continuation-in-part of Ser. No. 319,485, Mar. 6, 1989, abandoned.

[51] **Int. Cl.⁵** **A61K 37/22; C07H 19/06; C07H 19/10; C07H 19/16**[52] **U.S. Cl.** **424/450; 514/45; 514/49; 514/50; 514/51; 544/276; 544/277; 536/26.7; 536/26.8; 536/26.9; 536/26.23; 536/26.5; 536/26.71**[58] **Field of Search** **536/29, 27-28; 514/12, 808, 159, 49-51; 424/450, 38; 544/276, 277**[56] **References Cited****U.S. PATENT DOCUMENTS**

4,283,394	8/1981	West et al.	514/26
4,291,024	9/1981	Turcotte	514/48
4,471,113	9/1984	MacCoss	536/29
4,579,849	4/1986	MacCoss	514/262
4,622,392	11/1986	Hong et al.	536/29
4,670,424	6/1987	MacCoss	514/81
4,692,433	9/1987	Hostetler	514/12

FOREIGN PATENT DOCUMENTS

0056265	1/1981	European Pat. Off.
0122151	10/1984	European Pat. Off.
0199451	3/1986	European Pat. Off.
0254268	7/1987	European Pat. Off.
0286418	4/1988	European Pat. Off.
0311100	10/1988	European Pat. Off.
8807532	10/1988	European Pat. Off.
8902733	4/1989	World Int. Prop. O.

OTHER PUBLICATIONS

Richman, D. D., Kornbluth, R. S. and Carson, D. A., "Failure of Dideoxynucleosides to Inhibit Human Immunodeficiency Virus Replication in Cultured Human Macrophages", (1987), J. Exp. Med. 166: 1144-1149.
(List continued on next page.)

Primary Examiner—Johnnie R. Brown*Assistant Examiner*—Gary L. Kunz*Attorney, Agent, or Firm*—Knobbe, Martens, Olson & Bear[57] **ABSTRACT**

Compounds are disclosed for treating AIDS, herpes, and other viral infections by means of lipid derivatives of antiviral agents. The compounds consist of nucleoside analogues having antiviral activity which are linked, commonly through a phosphate group at the 5' position of the pentose residue, to one of a selected group of lipids. The lipophilic nature of these compounds provide advantages over the use of the nucleoside analogue alone. It also makes it possible to incorporate them into the lamellar structure of liposomes, either alone or combined with similar molecules. In the form of liposomes, these antiviral agents are preferentially taken up by macrophages and monocytes, cells which have been found to harbor the target HIV virus. Additional site specificity may be incorporated into the liposomes with the addition of ligands, such as monoclonal antibodies or other peptides or proteins which bind to viral proteins. Effective nucleoside analogues are dideoxynucleosides, azidothymine (AZT), and acyclovir; lipid groups may be glycolipids, sphingolipids, phospholipids or fatty acids. The compounds persist, after intracellular hydrolysis, as phosphorylated or non-phosphorylated antiviral nucleosides. The compounds are effective in improving the efficacy of antiviral nucleoside analogues by prolonging the antiviral activity after the administration of the drug has ended, and in preventing retroviral replication in HIV infections which have become resistant to therapy with conventional forms of the antiretroviral agents.

5 Claims, 3 Drawing Sheets